


LISTING OF THE CLAIMS

1-44 (Cancelled)

45. (Currently amended) Carboxamide compounds comprised of general formula I.1



wherein

U, V independently of one another denote CH ,


R²³, R²⁴ independently of one another denote H, F or methyl,

R²⁵, R²⁶, R²⁷ independently of one another have one of the meanings given for R²⁰ or in the case of a phenyl group also simply denote nitro, while residues R²⁵, R²⁶, R²⁷ occurring several times may have identical or different meanings, and


j is 0, 1, 2, 3 or 4 and

m, n independently of one another represent 0, 1 or 2.

R^1, R^2 independently of one another denote H, a C_{1-8} -alkyl or C_{3-7} -cycloalkyl group optionally substituted by the group R^{11} or a phenyl group optionally mono- or polysubstituted by the group R^{12} and/or monosubstituted by nitro, or



is defined according to one of the following partial formulae



wherein one or more H atoms of the heterocycle formed by the group R^1R^2N- may be replaced by R^{14} and the ring connected to the heterocycle formed by the group R^1R^2N- may be mono- or polysubstituted at one or more C atoms by R^{20} , and in the case of a phenyl ring may also

additionally be monosubstituted by nitro;

R³ denotes H, methyl or ethyl,

X denotes a single bond or a C₁₋₈-alkylene bridge wherein

one or two C atoms independently of one another may be substituted by a hydroxy, ω-hydroxy-C₁₋₃-alkyl-, ω-(C₁₋₃-alkoxy)-C₁₋₃-alkyl- and/or C₁₋₃-alkoxy group and/or in each case with one or two identical or different C₁₋₆-alkyl groups, and/or

Cy denotes a carbo- or heterocyclic group selected from one of the following meanings

- a saturated 3- to 7-membered carbocyclic group,
- an unsaturated 5- to 7-membered carbocyclic group,
- a phenyl group,
- a saturated 4- to 7-membered or unsaturated 5- to 7-membered heterocyclic group with an N, O or S atom as heteroatom,
- a saturated or unsaturated 5- to 7-membered heterocyclic group with two or more N atoms or with one or two N atoms and an O or S atom as heteroatoms,
- an aromatic heterocyclic 5- or 6-membered group with one or more identical or different heteroatoms selected from N, O and/or S,

~~while~~ wherein the ~~above-mentioned~~ 4, 5, 6 or 7-membered groups may be connected via two common, adjacent C atoms, fused with a phenyl or pyridine ring, and

wherein the ~~above-mentioned~~ 5-, 6- or 7-membered groups one or two non-adjacent -CH₂-groups may be replaced by a -CO-, -C(=CH₂)-, -(SO)- or -(SO₂)-group, and

wherein the ~~above-mentioned~~ saturated 6- or 7-membered groups may also be present as bridged ring systems with an imino, N-(C₁₋₄-alkyl)-imino, methylene, C₁₋₄-alkyl-methylene or di-(C₁₋₄-alkyl)-methylene bridge, and

wherein the ~~above-mentioned~~ cyclic groups may be mono- or polysubstituted at one or more C atoms with R²⁰, and in the case of a phenyl group also additionally monosubstituted by nitro, and/or substituted by R²¹ at one or more N atoms,

R¹¹ denotes R¹⁵-O-, R¹⁵-O-CO-, R¹⁶R¹⁷N-, R¹⁸R¹⁹N-CO- or Cy-,

R¹² has one of the meanings given for R²⁰,

R¹⁴ denotes halogen, C₁₋₆-alkyl, R¹⁵-O-, R¹⁵-O-CO-, R¹⁶R¹⁷N-, R¹⁸R¹⁹N-CO-, R¹⁵-O-C₁₋₃-alkyl-, R¹⁵-O-CO-C₁₋₃-alkyl-, R¹⁶R¹⁷N-C₁₋₃-alkyl-, R¹⁸R¹⁹N-CO-C₁₋₃-alkyl- or Cy-C₁₋₃-alkyl-,

R¹⁵ denotes H, C₁₋₄-alkyl, C₃₋₇-cycloalkyl, C₃₋₇-cycloalkyl-C₁₋₃-alkyl-, phenyl, phenyl-C₁₋₃-alkyl- or pyridinyl,

R¹⁶ denotes H, C₁₋₆-alkyl, C₃₋₇-cycloalkyl, C₃₋₇-cycloalkyl-C₁₋₃-alkyl-, C₄₋₇-cycloalkenyl, C₄₋₇-cycloalkenyl-C₁₋₃-alkyl-, ω-hydroxy-C₂₋₃-alkyl-, ω-(C₁₋₃-alkoxy)-C₂₋₃-alkyl-, amino-C₁₋₆-alkyl-, C₁₋₃-alkyl-amino-C₁₋₆-alkyl- or

di-(C₁₋₃-alkyl)-amino-C₁₋₆-alkyl-,

R¹⁷ has one of the meanings given for R¹⁶ or denotes phenyl, phenyl-C₁₋₃-alkyl-, pyridinyl, dioxolan-2-yl, C₁₋₃-alkylcarbonyl, hydroxycarbonyl-C₁₋₃-alkyl-, C₁₋₄-alkoxycarbonyl, C₁₋₃-alkylcarbonylamino-C₂₋₃-alkyl-, C₁₋₃-alkylsulphonyl- or C₁₋₃-alkylsulphonylamino-C₂₋₃-alkyl-,

R¹⁸, R¹⁹ independently of one another denote H or C₁₋₆-alkyl,

R²⁰ denotes halogen, hydroxy, cyano, C₁₋₄-alkyl, C₃₋₇-cycloalkyl, hydroxy-C₁₋₃-alkyl, R²²-C₁₋₃-alkyl- or has one of the meanings given for R²²,

R²¹ denotes C₁₋₃-alkyl, ω-hydroxy-C₂₋₃-alkyl-, phenyl, phenyl-C₁₋₃-alkyl-, C₁₋₃-alkyl-carbonyl, carboxy, C₁₋₄-alkoxy-carbonyl, C₁₋₃-alkylsulphonyl, phenylcarbonyl or phenyl-C₁₋₃-alkyl-carbonyl,

R²² denotes pyridinyl, phenyl, phenyl-C₁₋₃-alkoxy-, C₁₋₃-alkoxy, C₁₋₃-alkylthio, carboxy, H-CO-, C₁₋₃-alkylcarbonyl, C₁₋₄-alkoxycarbonyl, aminocarbonyl, C₁₋₃-alkylaminocarbonyl, di-(C₁₋₃-alkyl)-aminocarbonyl, C₁₋₃-alkylsulphonyl-, C₁₋₃-alkyl-sulphinyl-, C₁₋₃-alkyl-sulphonylamino-, amino, C₁₋₃-alkylamino-, di-(C₁₋₃-alkyl)-amino-, phenyl-C₁₋₃-alkylamino- or N-(C₁₋₃-alkyl)-phenyl-C₁₋₃-alkylamino-, acetylamino-, propionylamino-, phenylcarbonyl, phenylcarbonylamino-, phenylcarbonylmethylamino-, hydroxyalkylaminocarbonyl, (4-morpholinyl)carbonyl, (1-pyrrolidinyl)-carbonyl, (1-piperidinyl)carbonyl, (hexahydro-1-azepinyl)carbonyl, (4-methyl-1-piperazinyl)carbonyl, methylenedioxy-, aminocarbonylamino- or alkylaminocarbonylamino-,

while in the groups and residues X, R¹ to R³ and R¹¹, R¹² and R¹⁴ to R²² in each case one or more C atoms may be mono- or polysubstituted by F and/or in each case one

or two C atoms independently of one another may be monosubstituted by Cl or Br, and/or in each case one or more phenyl rings independently of one another additionally have one, two or three substituents selected from the group F, Cl, Br, I, C₁₋₄-alkyl, C₁₋₄-alkoxy, difluoromethyl, trifluoromethyl, hydroxy, amino, C₁₋₃-alkylamino-, di-(C₁₋₃-alkyl)-amino-, acetylamino-, aminocarbonyl, CN, difluoromethoxy, trifluoromethoxy, amino-C₁₋₃-alkyl-, C₁₋₃-alkylamino-C₁₋₃-alkyl- and di-(C₁₋₃-alkyl)-amino-C₁₋₃-alkyl- and/or may be monosubstituted by nitro, and

the tautomers, diastereomers, enantiomers, mixtures thereof and the salts thereof.

46. (Previously presented) The carboxamide compounds according to claim 45, wherein R¹, R² independently of one another denote H, C₁₋₆-alkyl, C₃₋₇-cycloalkyl, C₃₋₇-cycloalkyl-C₁₋₃-alkyl-, ω-hydroxy-C₂₋₃-alkyl, ω-(C₁₋₃-alkoxy)-C₂₋₃-alkyl-, C₁₋₄-alkoxy-carbonyl-C₁₋₃-alkyl, amino-C₂₋₄-alkyl-, C₁₋₃-alkyl-amino-C₂₋₄-alkyl- or di-(C₁₋₃-alkyl)-amino-C₂₋₄-alkyl-, phenyl or phenyl-C₁₋₃-alkyl-,

while in the above mentioned groups and residues one or more C atoms may be mono- or polysubstituted by F and/or one or two C atoms independently of one another may be monosubstituted by Cl or Br and

the phenyl group may be mono- or polysubstituted by the group R¹² defined in claim 45 and/or monosubstituted by nitro.

47. (Previously presented) The carboxamide compounds according to claim 45, wherein R¹, R² independently of one another denote C₁₋₄-alkyl, C₃₋₇-cycloalkyl, C₃₋₇-cycloalkyl-C₁₋₃-alkyl-, ω-hydroxy-C₂₋₃-alkyl-, ω-(C₁₋₃-alkoxy)-C₂₋₃-alkyl-, C₁₋₄-alkoxy-carbonyl-C₁₋₃-alkyl-, while one of the groups R¹, R² may also denote H.

48. (Previously presented) The carboxamide compounds according to claim 45, wherein R^1 , R^2 form with the N atom of the group R^1R^2N - a pyrrolidine, piperidine or 2,5-dihydro-1H-pyrrole ring, wherein one or more H atoms may be replaced by R^{14} and wherein R^{14} is defined as in claim 45.
49. (Previously presented) The carboxamide compounds according to claim 45, wherein R^{14} is selected from the group consisting of halogen, C_{1-4} -alkyl, hydroxy, C_{1-4} -alkoxy, C_{1-4} -alkoxy- C_{1-3} -alkyl-, hydroxy- C_{1-3} -alkyl, C_{1-4} -alkoxy-carbonyl, C_{1-4} -alkoxy-carbonyl- C_{1-3} -alkyl-, C_{1-4} -alkoxy-carbonylamino-, C_{1-4} -alkoxy-carbonylamino- C_{1-3} -alkyl-, amino, (C_{1-4} -alkyl)-amino-, di-(C_{1-4} -alkyl)-amino-.
50. (Previously presented) The carboxamide compounds according to claim 45, wherein X denotes a single bond or an alkylene bridge selected from methylene, 1,2-ethylene, 1,3-propylene and 1,4-butylene,
- wherein one or two C atoms independently of one another may be substituted with a hydroxy, ω -hydroxy- C_{1-3} -alkyl, ω -(C_{1-3} -alkoxy)- C_{1-3} -alkyl- and/or C_{1-3} -alkoxy group and/or in each case may be substituted with one or two identical or different C_{1-4} -alkyl groups, and
- in each case one or more C atoms may be mono- or polysubstituted by F and/or in each case one or two C atoms independently of one another may be monosubstituted by Cl or Br.
51. (Previously presented) The carboxamide compounds according to claim 45, wherein X is $-CH_2-$, $-CH(CH_3)-$ or $-C(CH_3)_2-$.
52. (Previously presented) The carboxamide compounds according to claim 45, wherein R^3 is H or methyl.

53. (Previously presented) The carboxamide compounds according to claim 45, wherein the group U denotes an N atom and the group V denotes a C atom.
54. (Previously presented) The carboxamide compounds according to claim 45, wherein the group U denotes a C atom and the group V denotes a N atom.
55. (Previously presented) The carboxamide compounds according to claim 45, wherein the two groups U and V each denote a C atom.
56. (Previously presented) Carboxamide compounds according to claim 45, wherein

R^{25} ,
 R^{26} , R^{27} independently of one another denote F, Cl, Br, I, OH, cyano, methyl, difluoromethyl, trifluoromethyl, ethyl, n-propyl, iso-propyl, methoxy, difluoromethoxy, trifluoromethoxy, ethoxy, n-propoxy or iso-propoxy, in the case of a substitution of a phenyl group they may also denote nitro, while repeatedly occurring groups R^{25} , R^{26} , R^{27} may have identical or different meanings, and

j is 0, 1 or 2, and

m, n independently of one another are 0 or 1.

57. (Previously presented) The carboxamide compounds according to claim 45, wherein the group R^{11} is selected from C_{1-6} -cycloalkyl, hydroxy, C_{1-4} -alkoxy, amino, C_{1-4} -alkyl-amino- and di- $(C_{1-4}$ -alkyl)-amino-.

58. (Previously presented) The carboxamide compounds according to claim 45, wherein the group R²⁰ is selected from halogen, hydroxy, cyano, C₁₋₄-alkyl, C₃₋₇-cycloalkyl and hydroxy-C₁₋₃-alkyl.
59. (Previously presented) The carboxamide compounds according to claim 45, wherein the group R²⁰ is selected from among fluorine, chlorine, bromine, CF₃, C₁₋₄-alkyl and C₁₋₄-alkoxy.
60. (Currently amended) Carboxamide compounds according to claim 45 selected from among the formulae:
- (1) — 4'-chloro-biphenyl-4-carboxylic acid [2-(4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
 - (2) — 4'-chloro-biphenyl-4-carboxylic acid-[2-(4-diethylaminomethyl-phenyl)-ethyl]-amide
 - (3) — 4'-chloro-biphenyl-4-carboxylic acid-[2-(4-piperidin-1-ylmethyl-phenyl)-ethyl]-amide
 - (4) — 4'-methoxy-biphenyl-4-carboxylic acid-[2-(4-diethylaminomethyl-phenyl)-ethyl]-amide
 - (5) — 4'-chloro-biphenyl-4-carboxylic acid-[2-(4-diethylaminomethyl-phenyl)-ethyl]-methyl-amide
 - (6) — 4'-chloro-biphenyl-4-carboxylic acid-[2-(4-pyrrolidin-1-ylmethyl-phenyl)-propyl]-amide
 - (7) — 4'-chloro-biphenyl-4-carboxylic acid-[2-(3-methoxy-4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
 - (8) — 4'-chloro-biphenyl-4-carboxylic acid-{2-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-ethyl}-amide

- (9) — 4'-chloro-biphenyl-4-carboxylic acid-[2-methyl-2-(4-pyrrolidin-1-ylmethyl-phenyl)-propyl]-amide
- (10) — ~~4-benzyl-N-[2-(4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-benzamide~~
- (11) — 4'-chloro-biphenyl-4-carboxylic acid-[2-(2-fluoro-4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- (12) — 4'-chloro-biphenyl-4-carboxylic acid [2-(3-bromo-4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- (13) — 4'-chloro-biphenyl-4-carboxylic acid [2-(3-methyl-4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- (14) — 4'-chloro-biphenyl-4-carboxylic acid {2-[4-(1-ethyl-piperidin-2-yl)-phenyl]-ethyl}-amide
- (15) — 4'-chloro-biphenyl-4-carboxylic acid{2-[4-(4-acetyl-piperazin-1-ylmethyl)-phenyl]-ethyl}-amide
- (16) — 4'-chloro-biphenyl-4-carboxylic acid (2-{4-[(diisopropylamino)-methyl]-phenyl}-ethyl)-amide
- (17) — 4'-chloro-biphenyl-4-carboxylic acid {2-[3-bromo-4-(2,5-dihydro-pyrrol-1-ylmethyl)-phenyl]-ethyl}-amide
- (18) — 4'-chloro-biphenyl-4-carboxylic acid{2-[4-(2-dimethylaminomethyl-pyrrolidin-1-ylmethyl)-phenyl]-ethyl}-amide
- (19) — 4'-chloro-biphenyl-4-carboxylic acid{2-[4-(3-dimethylamino-pyrrolidin-1-ylmethyl)-phenyl]-ethyl}-amide
- (20) — 4'-chloro-biphenyl-4-carboxylic acid [2-(2-bromo-4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- (21) — 4'-chloro-biphenyl-4-carboxylic acid [2-(6-pyrrolidin-1-

ylmethyl-pyridin-3-yl)-ethyl]-amide

- ~~(22)~~— 4'-chloro-biphenyl-4-carboxylic acid [2-(2-nitro-4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- ~~(23)~~— 2',4'-dichloro-biphenyl-4-carboxylic acid [2-(4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- ~~(24)~~— 4'-chloro-biphenyl-4-carboxylic acid{2-[4-(3-amino-pyrrolidin-1-ylmethyl)-phenyl]-ethyl}-amide
- ~~(25)~~— 4'-chloro-biphenyl-4-carboxylic acid{2-[4-(2-aminomethyl-pyrrolidin-1-ylmethyl)-phenyl]-ethyl}-amide
- ~~(26)~~— 4'-chloro-biphenyl-4-carboxylic acid [2-(5-pyrrolidin-1-ylmethyl-pyridin-2-yl)-ethyl]-amide
- ~~(27)~~— 4'-chloro-biphenyl-4-carboxylic acid [2-(3-ethyl-4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- ~~(28)~~— 4'-bromo-biphenyl-4-carboxylic acid {2-[4-(2,5-dihydro-pyrrol-1-ylmethyl)-phenyl]-ethyl}-amide
- ~~(29)~~— 4'-chloro-biphenyl-4-carboxylic acid [2-(2-methyl-4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- ~~(30)~~— 4'-bromo-3-fluoro-biphenyl-4-carboxylic acid {2-[3-bromo-4-(2,5-dihydro-pyrrol-1-ylmethyl)-phenyl]-ethyl}-amide
- ~~(31)~~— 4'-chloro-2-fluoro-biphenyl-4-carboxylic acid [2-(4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- ~~(32)~~— 4'-ethyl-biphenyl-4-carboxylic acid [2-(4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- ~~(33)~~— 4'-chloro-biphenyl-4-carboxylic acid {2-[4-(2-methyl-piperidin-1-ylmethyl)-phenyl]-ethyl}-amide
- ~~(34)~~— 4'-chloro-biphenyl-4-carboxylic acid {2-[4-(2-methyl-

pyrrolidin-1-ylmethyl)-phenyl]-ethyl}-amide

- (35) — 4'-chloro-biphenyl-4-carboxylic acid (2-{4-
[(cyclopropylmethyl-amino)-methyl]-phenyl}-ethyl)-amide
- (36) — ~~4'-chloro-biphenyl-4-carboxylic acid {2-[4-(3,4-dihydro-1H-isoquinolin-2-ylmethyl)-phenyl]-ethyl}-amide~~
- (37) — 4'-chloro-biphenyl-4-carboxylic acid [2-(4-{[(2-hydroxy-ethyl)-methyl-amino]-methyl}-phenyl)-ethyl]-amide
- (38) — 4'-chloro-biphenyl-4-carboxylic acid {2-[4-(2,6-dimethyl-piperidin-1-ylmethyl)-phenyl]-ethyl}-amide
- (39) — 4'-chloro-biphenyl-4-carboxylic acid [2-(4-azetidin-1-ylmethyl-phenyl)-ethyl]-amide
- (40) — 3,4'-dichloro-biphenyl-4-carboxylic acid [2-(4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- (41) — 4'-fluoro-biphenyl-4-carboxylic acid [2-(4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- (42) — 4'-chloro-3-fluoro-biphenyl-4-carboxylic acid [2-(4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- (43) — 2'-fluoro-4'-chloro-biphenyl-4-carboxylic acid [2-(4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- (44) — 5-(4-chloro-phenyl)-pyridine-2-carboxylic acid [2-(4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- (45) — 4'-chloro-biphenyl-4-carboxylic acid {2-[4-(2,5-dihydro-pyrrol-1-ylmethyl)-phenyl]-ethyl}-amide
- (46) — 4'-bromo-biphenyl-4-carboxylic acid [2-(4-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-amide
- (47) — 4'-chloro-biphenyl-4-carboxylic acid {2-[4-(1-pyrrolidin-1-yl-ethyl)-phenyl]-ethyl}-amide

and the pharmaceutically acceptable salts thereof.

61. (Previously presented) The physiologically acceptable salts of the carboxamide compounds according to claim 45.

62. (Cancelled)

63. (Previously presented) A composition comprised of at least one carboxamide compound according to claim 45 further comprised of one or more physiologically acceptable excipients.

64. (Previously presented) A pharmaceutical composition, comprising at least one carboxamide compound according to claim 45 optionally together with one or more inert carriers and/or diluents.

65. (Cancelled)

66. (Previously presented) A method of reducing the body weight and/or prevention of an increase of body weight of a mammal comprised of the administration of a therapeutically effective amount of a carboxamide compound according to claim 45 to a mammal or a pharmaceutically acceptable salt thereof.

67. (Previously presented) Use of at least one carboxamide compound according to claim 45 for preparing a pharmaceutical composition with an MCH-receptor-antagonistic activity.

68. (Cancelled)

69. (Currently amended) A method of treating ~~metabolic disorders and/or eating disorders~~, obesity, ~~bulimia, bulimia nervosa, cachexia, anorexia nervosa and hyperphagia~~ comprised of the administration to a patient in need thereof of a pharmaceutically effective amount of a carboxamide compound according to claim 45 or a pharmaceutically acceptable salt thereof.

70. (Currently amended) Method of preventing and/or treating diseases and/or disorders associated with obesity[[.]] selected from ~~particularly~~ diabetes, ~~especially~~ type II diabetes, complications of diabetes ~~including~~ selected from diabetic retinopathy, diabetic neuropathy, diabetic nephropathy, insulin resistance, pathological glucose tolerance, encephalorrhagia, cardiac insufficiency, cardiovascular diseases, ~~particularly~~ arteriosclerosis and high blood pressure, arthritis and gonitis, comprised of administering to a patient in need thereof a therapeutically effective amount of a compound according to claim 45 or a pharmaceutically acceptable salt thereof.

71. (Withdrawn) Method of preventing or treating hyperlipidaemia, cellulitis, fat accumulation, malignant mastocytosis, systemic mastocytosis, emotional disorders, affective disorders, depression, anxiety, sleep disorders, reproductive disorders, sexual disorders, memory disorders, epilepsy, forms of dementia and hormonal disorders comprised of administering to a patient in need thereof therapeutically effective amount of a carboxamide compound according to claim 45 or a pharmaceutically acceptable salt thereof.

72. (Withdrawn) A method of preventing and/or treating micturition disorders, selected from a list consisting of urinary incontinence, hyperactive urinary bladder, urgency, nycturia and enuresis comprised of administering to a patient in need thereof a therapeutically effective amount of a compound according to claim 45 or a pharmaceutically acceptable salt thereof.

73. (Withdrawn) Pharmaceutical compositions, comprised of a first active substance selected from the carboxamide compounds according to claim 1 and a second active substance selected from the group comprised of active substances for the treatment of diabetes, active substances for the treatment of diabetic complications, active substances for the treatment of obesity, other than MCH antagonists, active substances for the treatment of high blood pressure, active substances for the treatment of hyperlipidaemia, including arteriosclerosis, active substances for the treatment of arthritis, active substances for the treatment of anxiety states and active substances for the treatment of depression, optionally together with one or more inert carriers and/or diluents.